ELSEVIER

Gene Expression Patterns 1 (2002) 213–214

GENE EXPRESSION PATTERNS

www.elsevier.com/locate/bres

Author Index

Abramovitz, R., see Sharon, R. (1) 5
Ahi, J., see Nijholt, I. (1) 101
Aizawa, T., see Kuromitsu, J. (1) 17
Akiba, Y., Suzuki, R., Saito-Saino, S., Owada, Y., Sakagami, H., Watanabe, M. and Kondo, H.
Localization of mRNAs for phosphatidylinositol phosphate kinases in the mouse brain during development (1) 123
Alvarez-Bolado, G., see Moeller, C. (1) 199
Andersson, O. and von Euler, G.
Characterization and expression of the gene encoding membralin, an evolutionary conserved protein expressed in the central nervous system (1) 205
Bächner, D., Schröder, D. and Gross, G.
mRNA expression of the murine glycoprotein (transmembrane) nmb (*Gpnmb*) gene is linked to the developing retinal pigment epithelium and iris (1) 159
Barber, M.C., see Sorensen, A. (1) 167
Bird, P.I., see Kishi, T. (1) 175
Blank, T., see Nijholt, I. (1) 101
Carson, M.J., see Danielson, P.E. (1) 83
Chen, Z., Ge, B., Hudson, T.J. and Rozen, R.
Microarray analysis of brain RNA in mice with methylenetetrahydrofolate reductase deficiency and hyperhomocysteinemia (1) 89
Chiaramello, A., see Uittenbogaard, M. (1) 115
Chuai, M.-L., see Katsube, K.-i. (1) 61
Curran, T., see Magdaleno, S. (1) 143
Curran-Rauhut, M.A. and Petersen, S.L.
The distribution of progestin receptor mRNA in rat brainstem (1) 151
Danielson, P.E., Sautkulis, L.N., Foye, P.E., Hedlund, P.B. and Carson, M.J.
A novel mRNA expressed along brain ventricles (1) 83
Eichele, G., see Moeller, C. (1) 199
Feng, Y., Reznik, S.E. and Fricker, L.D.
ProSAAS and prohormone convertase 1 are broadly expressed during mouse development (1) 135
Foye, P.E., see Danielson, P.E. (1) 83
Fricker, L.D., see Feng, Y. (1) 135
Furuichi, T., see Kagami, Y. (1) 39
Ge, B., see Chen, Z. (1) 89
González, A., López, J.M. and Marín, O.
Expression pattern of the homeobox protein NKX2-1 in the developing *Xenopus* forebrain (1) 181
Gonzalez-Nicolini, V. and McGinty, J.F.
Gene expression profile from the striatum of amphetamine-treated rats: a cDNA array and in situ hybridization histochemical study (1) 193
Gross, G., see Bächner, D. (1) 159
Guan, X.-M., Yu, H., Jiang, Q., Van der Ploeg, L.H.T. and Liu, Q.
Distribution of neuromedin U receptor subtype 2 mRNA in the rat brain (1) 1
Haga, S., see Kuromitsu, J. (1) 17
Hashimoto, N., see Yamanaka, H. (1) 107
Hedlund, P.B., see Danielson, P.E. (1) 83
Hellebrand, S., Wittenberger, T., Schaller, H.C. and Hermans-Borgmeyer, I.
Gpr85, a novel member of the G-protein coupled receptor family, prominently expressed in the developing mouse cerebral cortex (1) 13
Hermans-Borgmeyer, I., see Hellebrand, S. (1) 13
Hudson, T.J., see Chen, Z. (1) 89
Ikeda, K., see Kuromitsu, J. (1) 17
Jiang, Q., see Guan, X.-M. (1) 1
Kabasawa, Y., see Katsube, K.-i. (1) 61
Kagami, Y. and Furuichi, T.
Investigation of differentially expressed genes during the development of mouse cerebellum (1) 39
Kato, K., see Kishi, T. (1) 175
Katsube, K.-i., Chuai, M.-L., Liu, Y.-C., Kabasawa, Y., Takagi, M., Perbal, B. and Sakamoto, K.
The expression of chicken *NOV*, a member of the CCN gene family, in early stage development (1) 61
Kawai, T., see Kuromitsu, J. (1) 17
Kishi, T., Matsuhashi, H., Bird, P.I. and Kato, K.
Distribution of serine proteinase inhibitor, clade B, member 6 (*Serpinb6*) in the adult mouse brain (1) 175
Kitamoto, T., see Salvaterra, P.M. (1) 73
Kondo, H., see Akiba, Y. (1) 123
Koyama, K., see Yamanaka, H. (1) 107
Kramer, P.R. and Wray, S.
Nasal embryonic LHRH factor (NELF) expression within the CNS and PNS of the rodent (1) 23
Kuromitsu, J., Yokoi, A., Kawai, T., Nagasu, T., Aizawa, T., Haga, S. and Ikeda, K.
Reduced neuropeptide Y mRNA levels in the frontal cortex of people with schizophrenia and bipolar disorder (1) 17
Kurschner, C., see Magdaleno, S. (1) 143
Lal, A., see Siu, I.-M. (1) 33
Liu, Q., see Guan, X.-M. (1) 1
Liu, Y.-C., see Katsube, K.-i. (1) 61
López, J.M., see González, A. (1) 181
Magdaleno, S., Northcutt, G.M., Curran, T. and Kurschner, C.
mPPP1R16B is a novel mouse protein phosphatase 1 targeting subunit whose mRNA is located in cell bodies and dendrites of neurons in four distinct regions of the brain (1) 143
Marín, O., see González, A. (1) 181
Matsuhashi, H., see Kishi, T. (1) 175
Maywood, E.S. and Mrosovsky, N.
A molecular explanation of interactions between photic and non-photic circadian clock-resetting stimuli (1) 27
Maywood, E.S. and Mrosovsky, N.
Erratum to: "A molecular explanation of interactions between photic and non-photic circadian clock-resetting stimuli". [Gene Expression Patterns 1 (2001) 27–31] (1) 141
McGinty, J.F., see Gonzalez-Nicolini, V. (1) 193
Miskin, R., see Sharon, R. (1) 5
Moeller, C., Yaylaoglu, M.B., Alvarez-Bolado, G., Thaller, C. and Eichele, G.
Murine *Lix1*, a novel marker for substantia nigra, cortical layer 5, and hindbrain structures (1) 199
Mrosovsky, N., see Maywood, E.S. (1) 27
Mrosovsky, N., see Maywood, E.S. (1) 141
Mueller, T., see Wullimann, M.F. (1) 187
Nagasu, T., see Kuromitsu, J. (1) 17
Nakagawa, H., see Yamanaka, H. (1) 107

1567-133X/02/$ – see front matter © 2002 Elsevier Science B.V. All rights reserved.
PII: S1567-133X(02)00044-3

Nakamura, Y., see Yamanaka, H. (1) 107
Nelson, P.A., Sutcliffe, J.G. and Thomas, E.A.
A new UDP-GalNAc:polypeptide *N*-acetylgalactosaminyltransferase mRNA exhibits predominant expression in the hypothalamus, thalamus and amygdala of mouse forebrain (1) 95
Nijholt, I., Blank, T., Ahi, J. and Spiess, J.
In vivo CREB phosphorylation mediated by dopamine and NMDA receptor activation in mouse hippocampus and caudate nucleus (1) 101
Noguchi, K., see Yamanaka, H. (1) 107
Northcutt, G.M., see Magdaleno, S. (1) 143
Owada, Y., see Akiba, Y. (1) 123
Perbal, B., see Katsube, K.-i. (1) 61
Petersen, S.L., see Curran-Rauhut, M.A. (1) 151
Price, N.T., see Sorensen, A. (1) 167
Reznik, S.E., see Feng, Y. (1) 135
Riggins, G.J., see Siu, I.-M. (1) 33
Rozen, R., see Chen, Z. (1) 89
Saito-Saino, S., see Akiba, Y. (1) 123
Sakagami, H., see Akiba, Y. (1) 123
Sakamoto, K., see Katsube, K.-i. (1) 61
Salvaterra, P.M. and Kitamoto, T.
Drosophila cholinergic neurons and processes visualized with Gal4/UAS–GFP (1) 73
Sautkulis, L.N., see Danielson, P.E. (1) 83
Schaller, H.C., see Hellebrand, S. (1) 13
Schröder, D., see Bächner, D. (1) 159
Sharon, R., Abramovitz, R. and Miskin, R.
Plasminogen activator inhibitor-2 (PAI-2) mRNA is localized in the accumbens nucleus of the mouse brain and is induced in specific brain sites after kainate excitation (1) 5
Siu, I.-M., Lal, A. and Riggins, G.J.
A database for regional gene expression in the human brain (1) 33
Sorensen, A., Travers, M.T., Vernon, R.G., Price, N.T. and Barber, M.C.
Localization of messenger RNAs encoding enzymes associated with malonyl-CoA metabolism in mouse brain (1) 167
Spiess, J., see Nijholt, I. (1) 101
Sutcliffe, J.G., see Nelson, P.A. (1) 95
Suzuki, R., see Akiba, Y. (1) 123
Takagi, M., see Katsube, K.-i. (1) 61
Thaller, C., see Moeller, C. (1) 199
Thomas, E.A., see Nelson, P.A. (1) 95
Travers, M.T., see Sorensen, A. (1) 167
Uittenbogaard, M. and Chiaramello, A.
Expression of the bHLH transcription factor *Tcf*12 (ME1) gene is linked to the expansion of precursor cell populations during neurogenesis (1) 115
Van der Ploeg, L.H.T., see Guan, X.-M. (1) 1
Vernon, R.G., see Sorensen, A. (1) 167
von Euler, G., see Andersson, O. (1) 205
Watanabe, M., see Akiba, Y. (1) 123
Wei, L.-N., see Zhou, F.C. (1) 67
Wittenberger, T., see Hellebrand, S. (1) 13
Wray, S., see Kramer, P.R. (1) 23
Wullimann, M.F. and Mueller, T.
Expression of *Zash-1a* in the postembryonic zebrafish brain allows comparison to mouse *Mash1* domains (1) 187
Yamanaka, H., Hashimoto, N., Koyama, K., Nakagawa, H., Nakamura, Y. and Noguchi, K.
Expression of Apc2 during mouse development (1) 107
Yaylaoglu, M.B., see Moeller, C. (1) 199
Yokoi, A., see Kuromitsu, J. (1) 17
Yu, H., see Guan, X.-M. (1) 1
Zhou, F.C. and Wei, L.-N.
Expression of cellular retinoic acid-binding protein I is specific to neurons in adult transgenic mouse brain (1) 67